

# Veyxyl® Tabs



## Tablets for cattle, pigs, horses, sheep and dogs

### NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Veyx-Pharma GmbH  
Soehreweg 6  
34639 Schwarzenborn  
Germany

### NAME OF THE VETERINARY MEDICINAL PRODUCT

Veyxyl® Tabs 800 mg/tablet  
Tablets for cattle, pigs, horses, sheep and dogs  
Amoxicillin as amoxicillin trihydrate

### STATEMENT OF THE ACTIVE SUBSTANCES AND OTHER INGREDIENTS

1 tablet contains:

*Active substance:*

800.0 mg Amoxicillin (equivalent to 918.48 mg Amoxicillin trihydrate)

### INDICATIONS

For treatment of the following diseases caused by gram-positive and/or gram-negative Amoxicillin-sensitive germs:

*Dogs; oral use:*

Infections of the lung and respiratory system, infections of the digestive tract, urogenital infections, localised infections, bacterial secondary infections following viral infections.

*Cattle, pigs, horses and sheep; intrauterine use:*

Infections of the endometrium, placental retention.

The sensitivity of the pathogens should be detected prior to the application of Veyxyl® Tabs. This is especially relevant for infections of the digestive tract due to very high rates of resistance to amoxicillin for *E. coli* and *Salmonellae*.

### CONTRAINDICATIONS

The treatment of animals hypersensitive to penicillins and cephalosporins; severe kidney function disorders with anuria and oliguria; the presence of  $\beta$ -lactamase-forming pathogens; oral application in ruminants and horses; oral application in rabbits, guinea-pigs, hamsters and other small rodents.  
Do not use Veyxyl® Tabs in animals weighing less than 40 kg.

### ADVERSE REACTIONS

Allergic reactions (allergic skin reactions, anaphylaxis)

In case of the occurrence of an allergic reaction the Veyxyl® Tabs treatment is to be stopped immediately.

Countermeasures:

In case of anaphylaxis: Adrenaline (epinephrine) and glucocorticoids i.v.

In case of allergic skin reactions: Antihistamines and/or glucocorticoids.

Following oral application, disorders of the gastrointestinal tract can occasionally occur (vomiting, diarrhoea, anorexia).

If you notice any adverse reaction not mentioned in this leaflet, please inform your veterinary surgeon or pharmacist.

### TARGET SPECIES

Cattle, pigs, horses, sheep, dogs

## DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

For oral use:

*Dogs above 40 kg body weight (BW):*

10 mg Amoxicillin per kg BW orally twice daily over 3 – 5 consecutive days (10 mg Amoxicillin per kg BW are equivalent to ½ tablet Veyxyl® Tabs per 40 kg BW). The tablet can also be given in the feed.

In case there is no significant improvement in the disease 3 days after the treatment commenced, then the diagnosis should be reviewed and, where appropriate, a change in therapy undertaken. Assessing the pathogens' sensitivity is recommended. The treatment should be maintained for 2 days after the clinical symptoms subside.

For intrauterine use:

*Dosage per treatment:*

*Horses:* 800 mg Amoxicillin, equivalent to 1 tablet Veyxyl® Tabs

*Cattle:* 400 - 800 mg Amoxicillin, equivalent to ½ – 1 tablet Veyxyl® Tabs

*Sheep, pigs:* 400 mg Amoxicillin, equivalent to ½ tablet Veyxyl® Tabs

Two treatments at an interval of two days. Assessing the pathogens' sensitivity is recommended.

## ADVICE ON CORRECT ADMINISTRATION

Not applicable.

## WITHDRAWAL PERIODS

<u>Cattle, sheep and horses:</u>	Meat and offal	4 days
	Milk	2 days
<u>Pigs:</u>	Meat and offal	4 days

## SPECIAL STORAGE PRECAUTIONS

Keep out of the reach and sight of children.

Do not use after the expiry date which is stated on the label and outer carton.

Store in a dry place not above +25 °C.

## SPECIAL WARNING(S)

Special warnings for each target species:

Not applicable.

Special precautions for use:

Special precautions for use in animals:

Not applicable.

*Special precautions to be taken by the person administering the veterinary medicinal product to animals:*

Persons sensitive to penicillins should avoid direct contact of the medicine with the skin or mucous membranes.

Use during pregnancy, lactation or lay:

Not applicable.

Interaction with other medicinal products and other forms of interaction:

In terms of the antibacterial effect, there exists a potential antagonism for penicillins and chemotherapeutics with a rapidly commencing bacteriostatic effect.

Overdose (symptoms, emergency procedures, antidotes), if necessary:

In case of overdosing symptoms of an excitation of the central nervous system and spasms/convulsions can occur. Veyxyl® Tabs treatment must be stopped immediately and an appropriate symptomatic treatment carried out (administration of benzodiazepines or barbiturates as antidotes).

Incompatibilities:

Not applicable.

## SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY

Any unused veterinary medicinal product or waste material derived from such veterinary medicinal product should be disposed of in accordance with local requirements. Ask your veterinary surgeon how to dispose of medicines no longer required.

## OTHER INFORMATION

Package sizes:

Slide box of 10 tablets

Refill package of 100 tablets

